

Flex Therapist CEUs

Suicide Prevention

1. Which screening tool is most commonly used in emergency departments to determine suicide risk?

- A. ASQ
 - B. Patient Safety Screener
 - C. Brief Suicide Safety Assessment
 - D. PHQ
-

2. What is the next step for a PT after identifying a patient is at high risk of suicide in a hospital setting?

- A. Contact RN and Mental Health Provider
 - B. Leave the Room for Safety
 - C. Mobilize Hospital Security
 - D. Mitigate Patient's Risk of Escaping
-

3. What group has the highest suicide rate in the state of Washington?

- A. Native American
 - B. Black
 - C. White
 - D. Asian
-

4. In what situation could a patient be involuntarily admitted for suicide risk?

- A. If a Patient is at Low Risk of Suicide
 - B. If a Family Member Asks
 - C. If the Evaluating Mental Health Provider Decides this
 - D. If the Patient Agrees to be Admitted
-

5. What is not a reason to call the National Suicide and Crisis Lifeline?

- A. Suicidal Ideation
 - B. Financial Worries
 - C. Advice on a Career Change
 - D. Emotional Distress
-

6. What is characterized by persistent thoughts of suicide?

- A. Suicidal Ideation**
 - B. Suicidal Attempt**
 - C. Suicidality**
 - D. Suicide**
-

7. What resource should a PT give a patient to search for counseling services anywhere in the United States?

- A. Mental Health Association**
 - B. Psychology Today**
 - C. Suicide Prevention Act**
 - D. NIH Mental Health Service Locator**
-

8. In which situation would a PT need to investigate into severe suicide risk?

- A. A Patient who Scored a 3 on the Suicide Behaviors Questionnaire**
 - B. A Patient with a Negative BSSA**
 - C. A Patient with a Score of 15 on the PHQ-9**
 - D. A Patient with a Score of 5 on the PHQ-9**
-

9. Why are elderly people represent high rates of suicide compared to other ages?

- A. Grief**
 - B. Lethal Means**
 - C. Financial Troubles**
 - D. All of These**
-

10. What should a PTA do in a hospital where a patient begins aggressively shouting at them and begins to walk toward them?

- A. Attempt to Calm the Patient**
 - B. Follow Protocol to Contact Hospital Security**
 - C. Walk Toward the Patient and Restrain Them**
 - D. Shout for Help**
-

11. What component is not part of a patient safety plan?

- A. Driving as a Self-Soothing Strategy**
 - B. Make a List of Triggering Events**
 - C. Clear a Room of Harmful Objects**
 - D. Contacting a Therapist in Suicidal Ideation**
-

12. What act is the most widespread effort in the United States to create funding to prevent self-harm?

- A. Suicide Act**
 - B. Suicide Prevention Act**
 - C. Suicide Cause Act**
 - D. None of These**
-

13. What disciplines of care should a PT know if a patient screens positive for moderate depression?

- A. Psychiatry**
 - B. Licensed Mental Health Counselor**
 - C. Primary Care**
 - D. Both A and B**
-

14. Which screening tool is best for identifying youth risk for death by suicide?

- A. ASQ**
 - B. PHQ**
 - C. BSSA**
 - D. Full Safety Assessment**
-

15. What resource should a PTA present to a patient who is at low risk of suicide?

- A. Emergency Safety Plan**
 - B. Suicide and Crisis Lifeline**
 - C. Intensive Outpatient Therapy Program**
 - D. Doctor referral**
-

16. Which risk factor for suicide is most directly linked to death by suicide?

- A. Mild Depression**
 - B. Anxiety**
 - C. Borderline Personality Disorder**
 - D. Loneliness**
-

17. What might a PT do if they overhear someone in a park (not their patient) talk about a suicide plan?

- A. Avoid Being Nosey and Walk Away**
- B. Call 911 and Report it**
- C. Approach the Person and Offer to Help**

D. Tell Another Person in the Park and then Leave

18. What factor explains why more men die by suicide even though more women attempt suicide?

- A. Lack of Resources**
 - B. Loneliness**
 - C. Higher Depression Rates**
 - D. More Lethal Suicide Means**
-

19. Which factor does not put a patient at a higher risk of suicide compared to the other factors?

- A. Past Suicide Attempt**
 - B. A Relationship Breakup**
 - C. Financial Hardship**
 - D. Homelessness**
-

20. In which group is suicide the most common cause of death in the world?

- A. Older Adults**
 - B. Hispanic Population**
 - C. Alaska Native Population**
 - D. Youth**
-

21. What should a PT do if an adult patient is hesitant to disclose information with their partner present for an outpatient PT examination?

- A. Nothing, the Partner Should Remain Present**
 - B. Ask if the Partner Would Go to the Waiting Room for a Few Minutes as Part of Evaluation Protocol**
 - C. Wait for an Opportunity to Ask Sensitive Questions until the Patient is Alone**
 - D. None of These**
-

22. Why might a person addicted to drugs be predisposed to suicide?

- A. Addiction Causes Isolation**
 - B. Addiction May Cause Financial Problems**
 - C. Addiction May Accompany Underlying Depression or Other Mental Health Diagnosis**
 - D. All of These**
-

23. What is the best strategy to reduce the number of suicide deaths in Washington and the remainder of the US?

- A. Early Access to Mental Health Services**
 - B. Psychiatric Inpatient Stay**
 - C. Suicide Screening of High-Risk Patients**
 - D. Dispersing the Suicide Crisis Line**
-

24. If a patient is resistant to participate in a suicide screening in a hospital, what should a PTA do?

- A. Skip The Suicide Screening**
 - B. Insist the Patient Participates**
 - C. Contact Mental Health Colleagues to Conduct the Screening**
 - D. Discuss the Patient's Refusal with PTA Colleagues**
-

25. What program might a patient discharging from a 24/7 inpatient psychiatric hospitalization undergo next as a transition before outpatient therapy?

- A. Post Psychiatric Hospitalization Therapy**
 - B. Partial Hospitalization**
 - C. Intensive Outpatient**
 - D. Either B or C**
-

26. What is appropriate language to introduce a suicide screening questionnaire?

- A. "This is a mandatory part of my evaluation."**
 - B. "Now I am going to ask a few sensitive questions I ask of all patients."**
 - C. "I need you to participate with a few additional questions."**
 - D. "Your history and risk factors point to a suicide attempt."**
-

27. What should a PT do if after alerting other providers in a hospital about active suicide risk, the patient becomes aggressive towards the PT?

- A. Follow Protocol for Mobilizing Hospital Security**
 - B. Try to Calm the Patient Down with Neutral Language**
 - C. Leave the Room**
 - D. Both A and C**
-

28. What is appropriate action for a school student who a PT finds to be at moderate risk of suicide?

- A. Contact School Mental Health Professionals**

- B. Refer the Student to Outpatient Mental Health**
 - C. Drive the Student to an Emergency Room**
 - D. Leave the Student Alone to Help Calm Him/Her**
-

29. Which provider will complete a full safety assessment for suicide risk?

- A. Physical Therapist**
 - B. Registered Nurse**
 - C. Licensed Mental Health Practitioner**
 - D. The Provider who Identified Suicide Risk**
-

30. What should a PTA do if their patient in home health discloses they have a plan to end their life?

- A. Call the Patient's Primary Care for a Referral**
 - B. Give the Patient the Suicide and Crisis Lifeline Number**
 - C. Call 911 and Stay with Patient**
 - D. Conduct a Full Safety Assessment**
-

Copyright © 2024 Flex Therapist CEUs

Visit us at <https://www.flextherapistceus.com>