

Flex Therapist CEUs

Achilles Tendon Injuries & Rehabilitation

1. Typically, how long is the Achilles tendon?

- A. 10 cm
 - B. 12 cm
 - C. 15 cm
 - D. 20 cm
-

2. What structures do the Achilles tendon attach to?

- A. Gastrocnemius and soleus
 - B. Plantaris and calcaneal tuberosity
 - C. Soleus and talocrual joint
 - D. Plantaris and lateral malleolus
-

3. What is a purpose action of the Achilles tendon?

- A. Propulsion of gait
 - B. Stablizing the heel
 - C. Ankle dorsiflexion
 - D. Ankle eversion
-

4. What are two differential diagnoses for suspect Achilles tendon pathology?

- A. Achilles tendon peritendinitis and meniscus tear
 - B. compartment syndrome and calf muscle strain
 - C. calcaneus fracture and 1st metatarsal fracture
 - D. DVT and Morton's neuroma
-

5. Where does insertional tendinopathy occur?

- A. Attachment from Achilles to calcaneus
 - B. Plantaris to calcaneus
 - C. Gastrocnemius to femur
 - D. Achilles tendon to gastrocnemius
-

6. What is another phrase for genu recurvatum?

- A. High arch
 - B. Knock kneed
 - C. Bow legged
 - D. Hyperextension
-

7. What is the normal range of motion in degrees for ankle dorsiflexion and plantarflexion, respectively?

- A. 30, 20
 - B. 20, 50
 - C. 50, 25
 - D. 20, 30
-

8. Which ankle joint contributes the most to ankle dorsiflexion and plantar flexion?

- A. Subtalar
 - B. Midtarsal
 - C. Talocrural
 - D. Tarsometatarsal
-

9. With Achilles tendinopathy rehabilitation, when should a patient progress to eccentric heel raises off of a step?

- A. At 2 weeks post injury
 - B. At one month post injury
 - C. At two months post injury
 - D. When pain is absent
-

10. Which ankle joint mobilization will increase dorsiflexion range of motion?

- A. Cuboid whip
 - B. Subtalar lateral glide
 - C. Subtalar medial glide
 - D. Talocrural joint thrust manipulation
-

11. Which intervention strategy is outdated and should not be used as a primary treatment for Achilles tendinopathy?

- A. Low level laser therapy
 - B. Iontophoresis
 - C. Eccentric loading
 - D. Activity modification
-

12. When should a patient typically progress to standing calf raises with Achilles partial tear?

- A. 3 weeks**
 - B. 7-12 weeks**
 - C. 12 weeks**
 - D. 5 weeks**
-

13. What age group most commonly fully ruptures the Achilles tendon?

- A. 2nd decade**
 - B. 5th or 6th decade**
 - C. 7th to 8th decade**
 - D. 3rd decade**
-

14. Which two tests are helpful in ruling in/out Achilles full rupture?

- A. Plantar flexion strength and Thompson's**
 - B. Thompson's and pain with rest**
 - C. Palpation of ruptured tendon and night pain**
 - D. Impaired plantar flexion strength and full active plantar flexion range of motion**
-

15. Which two imaging methods are used with Achilles pathology?

- A. MRI and PET scan**
 - B. Ultrasound and xray**
 - C. MRI and Ultrasound**
 - D. EMG and xray**
-

16. When can early range of motion typically begin after surgical repair of full rupture?

- A. 1 week**
 - B. 4 weeks**
 - C. 6 weeks**
 - D. 2 weeks**
-

17. How can an Achilles tendon rupture be managed?

- A. Rest and immobilization**
 - B. Nonoperative and operative**
 - C. Operative and immobilization for 8 weeks**
 - D. It will heal on its own**
-

18. What factors increase rerupture rate?

- A. Arterial disease, corticosteroid injection**
 - B. Active lifestyle**
 - C. Surgery and immobilization for 2 weeks**
 - D. Eccentric loading protocol**
-

19. What are two special tests for diagnosis of Achilles tendinopathy?

- A. Royal London Hospital test and Thompson test**
 - B. Hop test and DVT testing**
 - C. Hop test and painful arc test**
 - D. Soleus strength and ankle range of motion**
-

20. What rehabilitation intervention is accepted as most effective in returning function of Achilles tendon?

- A. Iontophoresis**
 - B. Stretching**
 - C. Manual therapy**
 - D. Eccentric loading protocol**
-

Copyright © 2024 Flex Therapist CEUs

Visit us at <https://www.flextherapistceus.com>