

Flex Therapist CEUs

Concussion Assessment Predictor Tool

1. The acute SCAT-3 symptom severity score is the most sensitive and significant predictor of recovery from sport-related concussions and hence of functional impairment and service utilization.

- A. True
 - B. False
-

2. All of the following are the primary causes of traumatic brain injury in patients seen in the Emergency Department, except for:

- A. Motor vehicle accidents
 - B. Sport-related injuries
 - C. Assaults
 - D. Falls
-

3. Findings from this study show that the presence and frequency of post-traumatic headache are associated with the SCAT-3 symptom severity score, especially with the _____ SCAT-3 symptoms.

- A. Somatic
 - B. Cognitive
 - C. Emotional
 - D. Sleep
-

4. The SCAT-3 symptom scores might be a useful tool for neurologists and headache specialists as they incorporate all of the following, except:

- A. Concentration
 - B. Memory
 - C. Balance
 - D. Coordination
-

5. TBI was found to be more prevalent in women than men in this study of non-athlete civilian patients.

- A. True
 - B. False
-

6. A prior history of headache and/or concussions are associated with the prevalence and intensity of posttraumatic headaches.

- A. True**
 - B. False**
-

7. Which of the following is associated with higher SCAT-3 symptom severity and SCAT-3 symptom scores?

- A. Post-traumatic headache prevalence**
 - B. Post-traumatic headache frequency**
 - C. Both post-traumatic headache prevalence and frequency are associated with higher SCAT-3 symptom severity and SCAT-3 symptom scores**
 - D. Neither post-traumatic headache prevalence nor frequency are associated with higher SCAT-3 symptom severity and SCAT-3 symptom scores**
-

8. Post-traumatic headache patients have higher SCAT-3 symptom scores than headache-free patients only when the “headache” symptom is counted when calculating the SCAT-3 symptom scores.

- A. True**
 - B. False**
-

9. Compared to the headache-free patients, when comparing SCAT-3 symptoms for each of the four symptom categories individually, post-traumatic headache patients have significantly higher cognitive SCAT-3 symptoms.

- A. True**
 - B. False**
-

10. When comparing the subjective and objective assessments, the SCAT-3 self-reported symptoms of balance, concentration, and memory correlate with the modified BESS and SAC scores, which supports the use of the scores among a heterogeneous population of concussion patients who are not necessarily athletes.

- A. True**
 - B. False**
-

11. A prior study on concussion assessment in an outpatient heterogeneous population showed that _____ is associated with worse SCAT-3 symptom scores.

- A. Older age**
- B. Male gender**

- C. Prior concussions**
 - D. Prior non-traumatic headache**
-

12. The SCAT-3 seems to be a good tool to evaluate post-traumatic headache in the general population of post-concussion patients.

- A. True**
 - B. False**
-

Copyright © 2024 Flex Therapist CEUs

Visit us at <https://www.flextherapistceus.com>