

# Flex Therapist CEUs

## Management of ACL Injuries in Children

1. How many main ligaments surround the knee to provide support?

- A. 4
  - B. 1
  - C. 3
  - D. 2
- 

2. What is not a risk factor for ACL tears?

- A. Improper footwear
  - B. Female gender
  - C. Sports
  - D. Hip weakness
- 

3. A major immediate sign of an ACL tear is;

- A. Nausea
  - B. Falling
  - C. A popping sound
  - D. Burning pain
- 

4. What is not a test used to determine a torn ACL?

- A. Reverse Clunk Test
  - B. Reverse Pivot Shift Test
  - C. Lachman's Sign Test
  - D. Clunk Test
- 

5. What can be used to show the extent of injury to the surrounding cartilage?

- A. X-ray
  - B. MRI
  - C. Anterior Drawer Test
  - D. CT scan
- 

6. What method is useful for alleviating pain and swelling throughout the process?

- A. SALT**
  - B. TIME**
  - C. POLICE**
  - D. FORT**
- 

**7. What movement is unlikely to contribute to an ACL injury?**

- A. Pivoting**
  - B. Cutting**
  - C. Jumping**
  - D. Squatting**
- 

**8. How long does it normally take to recover from ACL reconstruction?**

- A. 3-4 months**
  - B. 6-12 months**
  - C. 6-24 months**
  - D. 8-10 months**
- 

**9. Disturbance to the growth plate can result in leg length discrepancies and angular deformities.**

- A. True**
  - B. False**
- 

**10. Immediately after surgical reconstruction, the patient should begin exercise with resistance.**

- A. True**
  - B. False**
- 

**Copyright © 2024 Flex Therapist CEUs**

**Visit us at <https://www.flextherapistceus.com>**