

Flex Therapist CEUs

Rotator Cuff Tendinopathy and Upper Trapezius Effects

1. Rotator cuff tendinopathy is defined as a mechanical entrapment of the subacromial soft tissue underneath the acromial arch during arm elevation.

- A. True
 - B. False
-

2. How does the upper trapezius muscle contribute to normal scapular motion?

- A. The UT muscle elevates the scapula during arm elevation.
 - B. The UT muscle rotates the scapula during arm elevation.
 - C. The UT muscle elevates and rotates the scapula during arm elevation.
 - D. The UT muscle does not play a role in normal scapular motion.
-

3. Muscle shear modulus is linearly related to both active and passive muscle force.

- A. True
 - B. False
-

4. Compared to asymptomatic athletes, athletes with rotator cuff tendinopathy exhibited all of the following, except:

- A. Higher UT shear modulus when passively positioned at 30 degrees of shoulder abduction.
 - B. Higher UT shear modulus during the resting arm position at 0 degrees of shoulder abduction.
 - C. Higher UT shear modulus during active arm holding.
 - D. Compared to asymptomatic athletes, athletes with rotator cuff tendinopathy exhibited higher UT shear modulus under all arm positions.
-

5. The higher UT shear modulus values measured during the active tasks in athletes with rotator cuff tendinopathy are most likely explained by intrinsic changes in muscle mechanical properties.

- A. True
 - B. False
-

6. Increased UT activation has which of the following clinical consequences?

- A. Decreased activation level.
- B. A delayed onset activation of serratus anterior and lower trapezius.

- C. An altered scapular kinematics related to a greater superior translation of the scapula with less efficient upward rotation and posterior tipping.
- D. Increased UT activation has decreased activation level, a delayed onset activation of serratus anterior and lower trapezius, and an altered scapular kinematics related to a greater superior translation of the scapula with less efficient upward rotation and posterior tipping.
-

7. The difference in passive tension can only be assessed using:

- A. Electromyography
 - B. Elastography
 - C. Somatosensory evoked potentials
 - D. Autonomic testing
-

8. The upper trapezius is significantly more stretched at _____ degrees of shoulder abduction.

- A. 0
 - B. 30
 - C. 60
 - D. The UT experiences the same stretch at all degrees of shoulder abduction.
-

9. Active tension is related to the muscle extensibility.

- A. True
 - B. False
-

10. Individuals with forward shoulders may have stiffer upper trapezius.

- A. True
 - B. False
-

11. Athletes with UT passive shear modulus greater than _____ kPa may have higher risk of developing rotator cuff tendinopathy.

- A. 11.8
 - B. 12.0
 - C. 12.2
 - D. 12.4
-

12. This cross-sectional study was able to determine that the observed changes in UT shear modulus are the cause of rotator cuff tendinopathy.

- A. True
 - B. False
-

Copyright © 2024 Flex Therapist CEUs

Visit us at <https://www.flextherapistceus.com>